

EVALUATION OF SOME CONVOLUTION SUMS INVOLVING THE SUM OF DIVISORS FUNCTIONS

By

NATHALIE CHENG AND KENNETH S. WILLIAMS

(Received December 17, 2004; Revised June 30, 2005)

Abstract. The convolution sums $\sum_{m < n/2} \sigma_e(m)\sigma_f(n - 2m)$ and $\sum_{m < n/4} \sigma_e(m)\sigma_f(n - 4m)$ are evaluated explicitly for certain values of e and f and all positive integers n .

1. Introduction

For $e \in \mathbb{N}$ and $n \in \mathbb{N}$ we set

$$\sigma_e(n) = \sum_{d|n} d^e.$$

If $n \notin \mathbb{N}$ we set $\sigma_e(n) = 0$. We also write $\sigma(n)$ for $\sigma_1(n)$. We define the convolution sum $S_{e,f}(n)$ ($e, f, n \in \mathbb{N}$) by

$$(1.1) \quad S_{e,f}(n) := \sum_{m=1}^{n-1} \sigma_e(m)\sigma_f(n-m).$$

We note that

$$(1.2) \quad S_{e,f}(n) = S_{f,e}(n).$$

Ramanujan's tau function $\tau(n)$ ($n \in \mathbb{N}$) is defined by

$$(1.3) \quad \Delta(q) = q \prod_{n=1}^{\infty} (1 - q^n)^{24} = \sum_{n=1}^{\infty} \tau(n)q^n, \quad q \in \mathbb{C}, \quad |q| < 1.$$

The first twenty values of $\tau(n)$ are given in the following table.

2000 Mathematics Subject Classification: 11A25, 11E25, 11F11

Key words and phrases: sums of divisors functions, convolution sums, triangular numbers

* Research of the second author was supported by Natural Sciences and Engineering Research Council of Canada grant A-7233

N. CHENG AND K.S. WILLIAMS

n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$	n	$\tau(n)$
1	1	6	-6048	11	534612	16	987136
2	-24	7	-16744	12	-370944	17	-6905934
3	252	8	84480	13	-577738	18	2727432
4	-1472	9	-113643	14	401856	19	10661420
5	4830	10	-115920	15	1211760	20	-7109760

A table of the values of $\tau(n)$ can be found in [12]. Ramanujan [11] and Lahiri [6], [7] have shown that $S_{e,f}(n)$ can be expressed as a linear combination of

$$(1.4) \quad \sigma_j(n) \quad (j = 1, 3, \dots, e + f + 1), \quad n\sigma_{e+f-1}, \quad \tau(n)$$

with rational coefficients for those pairs $(e, f) \in \mathbb{N}^2$ satisfying

$$(1.5) \quad e + f \leq 12, \quad e \equiv f \equiv 1 \pmod{2}.$$

Specifically they proved

$$(1.6) \quad S_{1,1}(n) = \frac{5}{12}\sigma_3(n) + \frac{(1-6n)}{12}\sigma(n),$$

$$(1.7) \quad S_{1,3}(n) = \frac{7}{80}\sigma_5(n) + \frac{(1-3n)}{24}\sigma_3(n) - \frac{1}{240}\sigma(n),$$

$$(1.8) \quad S_{1,5}(n) = \frac{5}{126}\sigma_7(n) + \frac{(1-2n)}{24}\sigma_5(n) + \frac{1}{504}\sigma(n),$$

$$(1.9) \quad S_{3,3}(n) = \frac{1}{120}\sigma_7(n) - \frac{1}{120}\sigma_3(n),$$

$$(1.10) \quad S_{1,7}(n) = \frac{11}{480}\sigma_9(n) + \frac{(2-3n)}{48}\sigma_7(n) - \frac{1}{480}\sigma(n),$$

$$(1.11) \quad S_{3,5}(n) = \frac{11}{5040}\sigma_9(n) - \frac{1}{240}\sigma_5(n) + \frac{1}{504}\sigma_3(n),$$

$$(1.12) \quad S_{1,9}(n) = \frac{455}{30404}\sigma_{11}(n) + \frac{(5-6n)}{120}\sigma_9(n) + \frac{1}{264}\sigma(n) - \frac{36}{3455}\tau(n),$$

$$(1.13) \quad S_{3,7}(n) = \frac{91}{110560}\sigma_{11}(n) - \frac{1}{240}\sigma_7(n) - \frac{1}{480}\sigma_3(n) + \frac{15}{2764}\tau(n),$$

$$(1.14) \quad S_{5,5}(n) = \frac{65}{174132}\sigma_{11}(n) + \frac{1}{252}\sigma_5(n) - \frac{3}{691}\tau(n),$$

$$(1.15) \quad S_{1,11}(n) = \frac{691}{65520}\sigma_{13}(n) + \frac{(1-n)}{24}\sigma_{11}(n) - \frac{691}{65520}\sigma(n),$$

$$(1.16) \quad S_{3,9}(n) = \frac{1}{2640}\sigma_{13}(n) - \frac{1}{240}\sigma_9(n) + \frac{1}{264}\sigma_3(n),$$

$$(1.17) \quad S_{5,7}(n) = \frac{1}{10080}\sigma_{13}(n) + \frac{1}{504}\sigma_7(n) - \frac{1}{480}\sigma_5(n).$$

CONVOLUTION SUMS

In 1997, Melfi [8], [9] considered among others the convolution sums

$$\sum_{m < n/2} \sigma(m)\sigma(n-2m), \quad \sum_{m < n/2} \sigma(m)\sigma_3(n-2m), \quad \text{and} \quad \sum_{m < n/2} \sigma_3(m)\sigma(n-2m),$$

when n is odd, and proved that

$$\begin{aligned} \sum_{m < n/2} \sigma(m)\sigma(n-2m) &= \frac{1}{12}\sigma_3(n) + \frac{(1-3n)}{24}\sigma(n), \quad n \equiv 1 \pmod{2}, \\ \sum_{m < n/2} \sigma(m)\sigma_3(n-2m) &= \frac{1}{48}\sigma_5(n) + \frac{(2-3n)}{48}\sigma_3(n), \quad n \equiv 1 \pmod{2}, \\ \sum_{m < n/2} \sigma_3(m)\sigma(n-2m) &= \frac{1}{240}\sigma_5(n) - \frac{1}{240}\sigma(n), \quad n \equiv 1 \pmod{2}. \end{aligned}$$

For $e, f, n \in \mathbb{N}$ we set

$$(1.18) \quad T_{e,f}(n) = \sum_{m < n/2} \sigma_e(m)\sigma_f(n-2m)$$

and

$$(1.19) \quad U_{e,f}(n) = \sum_{m < n/4} \sigma_e(m)\sigma_f(n-4m).$$

In 2002 Huard, Ou, Spearman and Williams [4] evaluated $T_{e,f}(n)$ when $(e, f) = (1, 1), (1, 3), (3, 1)$ and $U_{e,f}(n)$ when $(e, f) = (1, 1)$ for all $n \in \mathbb{N}$, using only elementary methods. They proved

$$\begin{aligned} T_{1,1}(n) &= \frac{1}{12}\sigma_3(n) + \frac{1}{3}\sigma_3(n/2) + \frac{(1-3n)}{24}\sigma(n) + \frac{(1-6n)}{24}\sigma(n/2), \\ T_{1,3}(n) &= \frac{1}{48}\sigma_5(n) + \frac{1}{15}\sigma_5(n/2) + \frac{(2-3n)}{48}\sigma_3(n) - \frac{1}{240}\sigma(n/2), \\ T_{3,1}(n) &= \frac{1}{240}\sigma_5(n) + \frac{1}{12}\sigma_5(n/2) + \frac{(1-3n)}{24}\sigma_3(n/2) - \frac{1}{240}\sigma(n), \\ U_{1,1}(n) &= \frac{1}{48}\sigma_3(n) + \frac{1}{16}\sigma_3(n/2) + \frac{1}{3}\sigma_3(n/4) + \frac{(2-3n)}{48}\sigma(n) + \frac{(1-6n)}{24}\sigma(n/4). \end{aligned}$$

In this paper we evaluate $T_{e,f}(n)$ and $U_{e,f}(n)$ for all $n \in \mathbb{N}$ for those pairs $(e, f) \in \mathbb{N}^2$ satisfying $e + f \leq 10$, $e \equiv f \equiv 1 \pmod{2}$ (Sections 3-7). In Section 8 we use the values of $T_{3,1}(n)$ and $U_{3,1}(n)$ to reprove in a very simple way the formula of Ono, Robins and Wahl for the number of representations of $n \in \mathbb{N}$ as the sum of 12 triangular numbers [10, Theorem 7, p. 85].

2. Preliminary results

We begin by recalling Ramanujan's functions

$$(2.1) \quad L(q) = 1 - 24 \sum_{n=1}^{\infty} \sigma(n)q^n, \quad |q| < 1,$$

$$(2.2) \quad M(q) = 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n)q^n, \quad |q| < 1,$$

$$(2.3) \quad N(q) = 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n)q^n, \quad |q| < 1.$$

Ramanujan [11] and Lahiri [6], [7] actually use P , Q , R in place of L , M , N but we follow the usage of Berndt [1, p. 318]. For $k \in \mathbb{N}$ with $k > 1$ the Eisenstein series

$$E_{2k}(\tau) := 1 + \frac{2}{\zeta(1-2k)} \sum_{n=1}^{\infty} \sigma_{2k-1}(n)q^n,$$

where $\tau \in H = \{x + iy \in \mathbb{C} \mid y > 0\}$ and $q = e^{2\pi i\tau}$, is a modular form of weight $2k$ for the modular group $\Gamma = SL(2, \mathbb{Z})$. Here as usual $\zeta(s)$ denotes the Riemann zeta function. As $\zeta(-3) = 1/120$ and $\zeta(-5) = -1/252$ we have $E_4(\tau) = M(q)$ and $E_6(\tau) = N(q)$. The function $E_2(\tau) := 1 - 24 \sum_{n=1}^{\infty} \sigma(n)q^n = L(q)$ is not a modular form but is transformed under the action of Γ by

$$E_2\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^2 E_2(\tau) + \frac{6}{\pi i} c(c\tau + d), \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma,$$

see [1, p. 318]. We need the following identities which can be found in Lahiri [6], [7].

$$(2.4) \quad L^2(q) = 1 - 2^5 \cdot 3^2 \sum_{n=1}^{\infty} n\sigma(n)q^n + 2^4 \cdot 3 \cdot 5 \sum_{n=1}^{\infty} \sigma_3(n)q^n,$$

$$(2.5) \quad M^2(q) = 1 + 2^5 \cdot 3 \cdot 5 \sum_{n=1}^{\infty} \sigma_7(n)q^n,$$

$$(2.6) \quad M^3(q) = 1 + \frac{2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 13}{691} \sum_{n=1}^{\infty} \sigma_{11}(n)q^n + \frac{2^7 \cdot 3^3 \cdot 5^3}{691} \sum_{n=1}^{\infty} \tau(n)q^n,$$

$$(2.7) \quad N^2(q) = 1 + \frac{2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 13}{691} \sum_{n=1}^{\infty} \sigma_{11}(n)q^n - \frac{2^6 \cdot 3^5 \cdot 7^2}{691} \sum_{n=1}^{\infty} \tau(n)q^n,$$

CONVOLUTION SUMS

$$(2.8) \quad L(q)M(q) = 1 + 2^4 \cdot 3^2 \cdot 5 \sum_{n=1}^{\infty} n\sigma_3(n)q^n - 2^3 \cdot 3^2 \cdot 7 \sum_{n=1}^{\infty} \sigma_5(n)q^n,$$

$$(2.9) \quad L(q)M^2(q) = 1 + 2^4 \cdot 3^2 \cdot 5 \sum_{n=1}^{\infty} n\sigma_7(n)q^n - 2^3 \cdot 3 \cdot 11 \sum_{n=1}^{\infty} \sigma_9(n)q^n,$$

$$(2.10) \quad L(q)N(q) = 1 - 2^4 \cdot 3^2 \cdot 7 \sum_{n=1}^{\infty} n\sigma_5(n)q^n + 2^5 \cdot 3 \cdot 5 \sum_{n=1}^{\infty} \sigma_7(n)q^n,$$

$$(2.11) \quad M(q)N(q) = 1 - 2^3 \cdot 3 \cdot 11 \sum_{n=1}^{\infty} \sigma_9(n)q^n,$$

$$(2.12) \quad L(q)M(q)N(q) = 1 - \frac{2^4 \cdot 3^2 \cdot 11}{5} \sum_{n=1}^{\infty} n\sigma_9(n)q^n \\ + \frac{2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 13}{691} \sum_{n=1}^{\infty} \sigma_{11}(n)q^n - \frac{2^3 \cdot 3^4 \cdot 11}{5 \cdot 691} \sum_{n=1}^{\infty} \tau(n)q^n.$$

From this point on we restrict q to be a real number satisfying $0 < q < 1$. Then $0 < -\log q < \infty$. As usual we denote the Gaussian hypergeometric function by ${}_2F_1(a, b; c; x)$. The derivative y' of the function

$$y = \frac{\pi {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; 1-x\right)}{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right)}$$

is given by

$$y' = -\frac{x^{-1}(1-x)^{-1}}{\left({}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right)\right)^2},$$

see [1, p. 87]. For $0 < x < 1$, we have

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right) = \sum_{n=0}^{\infty} \frac{1}{2^{4n}} \binom{2n}{n}^2 x^n > 0$$

so that $y' < 0$ for $0 < x < 1$. Hence y is a decreasing function of x for $0 < x < 1$. As $y(0) = \infty$ and $y(1) = 0$, the function y decreases from ∞ to 0 as x increases from 0 to 1. Hence there is a unique value of x between 0 and 1 such that

$$(2.13) \quad y = \frac{\pi {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; 1-x\right)}{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right)} = -\log q.$$

Thus

$$(2.14) \quad q = e^{-y} = e^{-\frac{\pi {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; 1-x\right)}{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right)}}$$

We also set

$$(2.15) \quad w = {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right).$$

The following formulae are proved in [2, pp. 126–129]

$$(2.16) \quad L(q) = (1 - 5x)w^2 + 12x(1 - x)w \frac{dw}{dx},$$

$$(2.17) \quad M(q) = (1 + 14x + x^2)w^4,$$

$$(2.18) \quad N(q) = (1 + x)(1 - 34x + x^2)w^6,$$

$$(2.19) \quad L(q^2) = (1 - 2x)w^2 + 6x(1 - x)w \frac{dw}{dx},$$

$$(2.20) \quad M(q^2) = (1 - x + x^2)w^4,$$

$$(2.21) \quad N(q^2) = (1 + x)\left(1 - \frac{1}{2}x\right)(1 - 2x)w^6,$$

$$(2.22) \quad L(q^4) = \left(1 - \frac{5}{4}x\right)w^2 + 3x(1 - x)w \frac{dw}{dx},$$

$$(2.23) \quad M(q^4) = \left(1 - x + \frac{1}{16}x^2\right)w^4,$$

$$(2.24) \quad N(q^4) = \left(1 - \frac{1}{2}x\right)\left(1 - x - \frac{1}{32}x^2\right)w^6.$$

Finally, from (2.6) and (2.7), we obtain

$$(2.25) \quad M^3(q) - N^2(q) = 2^6 \cdot 3^3 \sum_{n=1}^{\infty} \tau(n)q^n.$$

Appealing to (2.17), (2.18) and (2.25), we deduce

$$(2.26) \quad \Delta(q) = \sum_{n=1}^{\infty} \tau(n)q^n = \frac{x(1-x)^4 w^{12}}{2^4}$$

Applying the principle of duplication [2, p. 125]

$$q \rightarrow q^2, \quad x \rightarrow \left(\frac{1 - \sqrt{1-x}}{1 + \sqrt{1-x}}\right)^2, \quad w \rightarrow w \left(\frac{1 + \sqrt{1-x}}{2}\right)$$

to (2.26), we obtain

$$(2.27) \quad \Delta(q^2) = \sum_{n=1}^{\infty} \tau(n)q^{2n} = \frac{x^2(1-x)^2 w^{12}}{2^8}$$

Applying the principle of duplication to (2.27), we obtain

$$(2.28) \quad \Delta(q^4) = \sum_{n=1}^{\infty} \tau(n)q^{4n} = \frac{x^4(1-x)w^{12}}{2^{16}}$$

3. Evaluation of $T_{e,f}(n)$ and $U_{e,f}(n)$ for $e + f = 2$

We begin by determining $L(q)L(q^2)$ and $L(q)L(q^4)$.

THEOREM 3.1.

(i) $L(q)L(q^2) = \frac{1}{4}L^2(q) + L^2(q^2) - \frac{1}{20}M(q) - \frac{1}{5}M(q^2),$

(ii) $L(q)L(q^4) = \frac{1}{8}L(q)^2 + 2L(q^4)^2 - \frac{3}{40}M(q) + \frac{3}{20}M(q^2) - \frac{6}{5}M(q^4).$

Proof. From (2.16) and (2.19) we obtain

(3.1) $-L(q) + 2L(q^2) = (1 + x)w^2.$

Squaring we deduce that

$$L^2(q) - 4L(q)L(q^2) + 4L^2(q^2) = (1 + 2x + x^2)w^4.$$

From (2.17) and (2.20) we have

$$M(q) + 4M(q^2) = (5 + 10x + 5x^2)w^4.$$

Hence

$$L^2(q) - 4L(q)L(q^2) + 4L^2(q^2) = \frac{1}{5}M(q) + \frac{4}{5}M(q^2),$$

from which part (i) follows. Part (ii) follows similarly. \square

From Theorem 3.1 we deduce the values of $T_{1,1}(n)$ and $U_{1,1}(n)$ mentioned at the end of Section 1. \blacktriangle

THEOREM 3.2.

(i) $T_{1,1}(n) = \frac{1}{12}\sigma_3(n) + \frac{1}{3}\sigma_3(n/2) + \frac{(1-3n)}{24}\sigma(n) + \frac{(1-6n)}{24}\sigma(n/2),$

(ii) $U_{1,1}(n) = \frac{1}{48}\sigma_3(n) + \frac{1}{16}\sigma_3(n/2) + \frac{1}{3}\sigma_3(n/4) + \frac{(2-3n)}{48}\sigma(n) + \frac{(1-6n)}{24}\sigma(n/4).$

Proof. We have

$$\begin{aligned} \sum_{n=1}^{\infty} T_{1,1}(n)q^n &= \sum_{n=1}^{\infty} \left(\sum_{m < n/2} \sigma(m)\sigma(n-2m) \right) q^n \\ &= \sum_{l=1}^{\infty} \sigma(l)q^l \sum_{m=1}^{\infty} \sigma(m)q^{2m} = \left(\frac{1-L(q)}{24} \right) \left(\frac{1-L(q^2)}{24} \right) \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2^6 \cdot 3^2} (1 - L(q) - L(q^2) + L(q)L(q^2)) \\
 &= \frac{1}{2^8 \cdot 3^2 \cdot 5} (20 - 20L(q) - 20L(q^2) + 5L^2(q) + 20L^2(q^2) \\
 &\quad - M(q) - 4M(q^2)),
 \end{aligned}$$

by Theorem 3.1 (i). Appealing to (2.1), (2.2) and (2.4), and equating the coefficients of q^n , we obtain the asserted formula for $T_{1,1}(n)$. The formula for $U_{1,1}(n)$ follows similarly from Theorem 3.1(ii), (2.1), (2.2) and (2.4). \square

Theorem 3.2 (i) is [4, Theorem 2, p. 247] and Theorem 3.2 (ii) is [4, Theorem 4, p. 249].

4. Evaluation of $T_{e,f}(n)$ and $U_{e,f}(n)$ for $e + f = 4$

We first determine $L(q)M(q^2)$, $L(q^2)M(q)$, $L(q)M(q^4)$ and $L(q^4)M(q)$. In order to do this we introduce the constants $a(n)$ ($n = 1, 2, \dots$) defined by

$$(4.1) \quad A(q) = \sum_{n=1}^{\infty} a(n)q^n = \Delta(q^2)^{1/2} = \frac{x(1-x)w^6}{2^4}.$$

The first twenty values of $a(n)$ are given in the following table.

n	$a(n)$	n	$a(n)$	n	$a(n)$	n	$a(n)$
1	1	6	0	11	540	16	0
2	0	7	-88	12	0	17	594
3	-12	8	0	13	-418	18	0
4	0	9	-99	14	0	19	836
5	54	10	0	15	-648	20	0

Similarly to the proof of Theorem 3.1 we obtain

THEOREM 4.1.

- (i) $L(q)M(q^2) = 2L(q^2)M(q^2) + \frac{1}{21}N(q) - \frac{22}{21}N(q^2)$,
- (ii) $M(q)L(q^2) = \frac{1}{2}L(q)M(q) - \frac{11}{42}N(q) + \frac{16}{21}N(q^2)$,
- (iii) $L(q)M(q^4) = 4L(q^4)M(q^4) + \frac{1}{336}N(q) + \frac{5}{112}N(q^2) - \frac{64}{21}N(q^4) - \frac{45}{2}A(q)$,
- (iv) $M(q)L(q^4) = \frac{1}{4}L(q)M(q) - \frac{4}{21}N(q) + \frac{5}{28}N(q^2) + \frac{16}{21}N(q^4) + 90A(q)$.

Appealing to (2.1), (2.2), (2.3), (2.8), (4.1) and Theorem 4.1, we obtain

CONVOLUTION SUMS

THEOREM 4.2.

$$\begin{aligned}
 \text{(i)} \quad T_{3,1}(n) &= \frac{1}{240}\sigma_5(n) + \frac{1}{12}\sigma_5(n/2) + \frac{(1-3n)}{24}\sigma_3(n/2) - \frac{1}{240}\sigma(n), \\
 \text{(ii)} \quad T_{1,3}(n) &= \frac{1}{48}\sigma_5(n) + \frac{1}{15}\sigma_5(n/2) + \frac{(2-3n)}{48}\sigma_3(n) - \frac{1}{240}\sigma(n/2), \\
 \text{(iii)} \quad U_{3,1}(n) &= \frac{1}{3840}\sigma_5(n) + \frac{1}{256}\sigma_5(n/2) + \frac{1}{12}\sigma_5(n/4) + \frac{(1-3n)}{24}\sigma_3(n/4) \\
 &\quad - \frac{1}{240}\sigma(n) + \frac{1}{256}a(n), \\
 \text{(iv)} \quad U_{1,3}(n) &= \frac{1}{192}\sigma_5(n) + \frac{1}{64}\sigma_5(n/2) + \frac{1}{15}\sigma_5(n/4) + \frac{(4-3n)}{96}\sigma_3(n) \\
 &\quad - \frac{1}{240}\sigma(n/4) - \frac{1}{64}a(n).
 \end{aligned}$$

The values of $T_{1,3}(n)$ and $T_{3,1}(n)$ were given at the end of Section 1. Theorem 4.2 (i), (ii) comprise [4, Theorem 6, p. 250].

5. Evaluation of $T_{e,f}(n)$ and $U_{e,f}(n)$ for $e + f = 6$

We first determine $L(q)N(q^2)$, $M(q)M(q^2)$, $L(q^2)N(q)$, $L(q)N(q^4)$, $M(q)M(q^4)$ and $L(q^4)N(q)$. In order to do this we introduce the constants $b(n)$ ($n \in \mathbb{N}$) defined by

$$(5.1) \quad B(q) = \sum_{n=1}^{\infty} b(n)q^n = (\Delta(q)\Delta(q^2))^{1/3} = \frac{x(1-x)^2w^8}{2^4}$$

Hence

$$(5.2) \quad B(q^2) = \sum_{n=1}^{\infty} b(n)q^{2n} = \sum_{n=1}^{\infty} b(n/2)q^n = (\Delta(q^2)\Delta(q^4))^{1/3} = \frac{x^2(1-x)w^8}{2^8}$$

The first twenty values of $b(n)$ are given in the following table.

n	$b(n)$	n	$b(n)$	n	$b(n)$	n	$b(n)$
1	1	6	-96	11	1092	16	4096
2	-8	7	1016	12	768	17	14706
3	12	8	-512	13	1382	18	16344
4	64	9	-2043	14	-8128	19	-39940
5	-210	10	1680	15	-2520	20	-13440

Similarly to the proof of Theorem 3.1, we obtain the following result.

THEOREM 5.1.

- (i) $L(q)N(q^2) = 2L(q^2)N(q^2) + \frac{1}{85}M^2(q) - \frac{86}{85}M^2(q^2) - \frac{504}{17}B(q),$
- (ii) $M(q)M(q^2) = \frac{1}{17}M^2(q) + \frac{16}{17}M^2(q^2) + \frac{3600}{17}B(q),$
- (iii) $N(q)L(q^2) = \frac{1}{2}L(q)N(q) - \frac{43}{170}M^2(q) + \frac{64}{85}M^2(q^2) - \frac{2016}{17}B(q),$
- (iv) $L(q)N(q^4) = 4L(q^4)N(q^4) + \frac{1}{5440}M^2(q) + \frac{63}{5440}M^2(q^2) - \frac{256}{85}M^2(q^4)$
 $- \frac{819}{34}B(q) - \frac{4788}{17}B(q^2),$
- (v) $M(q)M(q^4) = \frac{1}{272}M^2(q) + \frac{15}{272}M^2(q^2) + \frac{16}{17}M^2(q^4) + \frac{4050}{17}B(q) + \frac{64800}{17}B(q^2),$
- (vi) $N(q)L(q^4) = \frac{1}{4}L(q)N(q) - \frac{16}{85}M^2(q) + \frac{63}{340}M^2(q^2) + \frac{64}{85}M^2(q^4)$
 $- \frac{4788}{17}B(q) - \frac{104832}{17}B(q^2).$

Appealing to (2.1), (2.2), (2.3), (2.5), (2.10), (5.1), (5.2) and Theorem 5.1, we obtain the next result.

THEOREM 5.2.

- (i) $T_{5,1}(n) = \frac{1}{2142}\sigma_7(n) + \frac{2}{51}\sigma_7(n/2) + \frac{(1-2n)}{24}\sigma_5(n/2) + \frac{1}{504}\sigma(n) - \frac{1}{408}b(n),$
- (ii) $T_{3,3}(n) = \frac{1}{2040}\sigma_7(n) + \frac{2}{255}\sigma_7(n/2) - \frac{1}{240}\sigma_3(n) - \frac{1}{240}\sigma_3(n/2) + \frac{1}{272}b(n),$
- (iii) $T_{1,5}(n) = \frac{1}{102}\sigma_7(n) + \frac{32}{1071}\sigma_7(n/2) + \frac{(1-n)}{24}\sigma_5(n) + \frac{1}{504}\sigma(n/2) - \frac{1}{102}b(n),$
- (iv) $U_{5,1}(n) = \frac{1}{137088}\sigma_7(n) + \frac{1}{2176}\sigma_7(n/2) + \frac{2}{51}\sigma_7(n/4) + \frac{(1-2n)}{24}\sigma_5(n/4)$
 $+ \frac{1}{504}\sigma(n) - \frac{13}{6528}b(n) - \frac{19}{816}b(n/2),$
- (v) $U_{3,3}(n) = \frac{1}{32640}\sigma_7(n) + \frac{1}{2176}\sigma_7(n/2) + \frac{2}{255}\sigma_7(n/4) - \frac{1}{240}\sigma_3(n) - \frac{1}{240}\sigma_3(n/4)$
 $+ \frac{9}{2176}b(n) + \frac{9}{136}b(n/2),$
- (vi) $U_{1,5}(n) = \frac{1}{408}\sigma_7(n) + \frac{1}{136}\sigma_7(n/2) + \frac{32}{1071}\sigma_7(n/4) + \frac{(2-n)}{48}\sigma_5(n) + \frac{1}{504}\sigma(n/4)$
 $- \frac{19}{816}b(n) - \frac{26}{51}b(n/2).$

The values of $3T_{1,5}(n) + 8T_{3,3}(n)$ and $2T_{3,3}(n) + 3T_{5,1}(n)$ have been derived in an elementary manner in [4, Theorem 15, p. 271]. We also note that [5, Theorem 8, p. 162] follows from Theorem 5.2.

In [5, Theorem 6, p. 157] it is shown that the number of representations of $n (\in \mathbb{N})$ as the sum of sixteen squares is given by

$$r_{16}(n) = \frac{32}{15}(-1)^n\sigma_7(n) + \frac{8192}{15}\sigma_7(n/2) + \frac{512}{15}(-1)^{n-1}\sigma_3(n) - \frac{512}{15}\sigma_3(n/2) + 8192(-1)^{n-1}T_{3,3}(n),$$

and, appealing to Theorem 5.2 (ii) for the value of $T_{3,3}(n)$, we obtain

$$r_{16}(n) = \frac{32}{17}(-1)^{n-1}(\sigma_7(n) - 256\sigma_7(n/2) + 16b(n)),$$

which is a result due to Glaisher [3, p. 480].

6. Evaluation of $T_{e,f}(n)$ and $U_{e,f}(n)$ for $e + f = 8$

We first determine $L(q)M^2(q^2)$, $M(q)N(q^2)$, $N(q)M(q^2)$, $M^2(q)L(q^2)$, $L(q)M^2(q^4)$, $M(q)N(q^4)$, $N(q)M(q^4)$ and $M^2(q)L(q^4)$. In order to do this we introduce the constants $c(n)$, $d(n)$ and $e(n)$ ($n \in \mathbb{N}$) defined by

$$(6.1) \quad C(q) = \sum_{n=1}^{\infty} c(n)q^n = (\Delta(q)^4\Delta(q^2))^{1/6} = \frac{x(1-x)^3w^{10}}{2^4},$$

$$(6.2) \quad D(q) = \sum_{n=1}^{\infty} d(n)q^n = (\Delta(q)^2\Delta(q^2)\Delta(q^4)^2)^{1/6} = \frac{x^2(1-x)^2w^{10}}{2^8}$$

$$(6.3) \quad E(q) = \sum_{n=1}^{\infty} e(n)q^n = (\Delta(q^2)\Delta(q^4)^4)^{1/6} = \frac{x^3(1-x)w^{10}}{2^{12}}$$

We observe that

$$(6.4) \quad C(q)E(q) = D(q)^2$$

so that

$$(6.5) \quad \sum_{m=1}^{n-1} c(m)e(n-m) = \sum_{m=1}^{n-1} d(m)d(n-m).$$

The first twenty values of $c(n)$, $d(n)$ and $e(n)$ are given in the following tables.

n	$c(n)$	n	$c(n)$	n	$c(n)$	n	$c(n)$
1	1	6	2496	11	-38996	16	-65536
2	-16	7	-4536	12	39936	17	311442
3	100	8	-4096	13	37806	18	-74448
4	-256	9	23085	14	15232	19	128244
5	-154	10	-13920	15	-146472	20	-222720

n	$d(n)$	n	$d(n)$	n	$d(n)$	n	$d(n)$
1	0	6	-156	11	-536	16	4096
2	1	7	112	12	-2496	17	-17472
3	-8	8	256	13	4384	18	4653
4	16	9	-576	14	-952	19	5848
5	32	10	870	15	336	20	13920

n	$e(n)$	n	$e(n)$	n	$e(n)$	n	$e(n)$
1	0	6	0	11	67	16	0
2	0	7	-14	12	0	17	2184
3	1	8	0	13	-548	18	0
4	0	9	72	14	0	19	-731
5	-4	10	0	15	-42	20	0

Similarly to the proof of Theorem 3.1, we obtain the following result.

THEOREM 6.1.

- (i) $L(q)M^2(q^2) = 2L(q^2)M^2(q^2) + \frac{1}{341}M(q)N(q) - \frac{342}{341}M(q^2)N(q^2) - \frac{720}{31}C(q) - \frac{23040}{31}D(q),$
- (ii) $M(q)N(q^2) = \frac{5}{341}M(q)N(q) + \frac{336}{341}M(q^2)N(q^2) + \frac{7560}{31}C(q) + \frac{241920}{31}D(q),$
- (iii) $N(q)M(q^2) = \frac{21}{341}M(q)N(q) + \frac{320}{341}M(q^2)N(q^2) - \frac{15120}{31}C(q) - \frac{483840}{31}D(q),$
- (iv) $M^2(q)L(q^2) = \frac{1}{2}L(q)M^2(q) - \frac{171}{682}M(q)N(q) + \frac{256}{341}M(q^2)N(q^2) + \frac{5760}{31}C(q) + \frac{184320}{31}D(q),$
- (v) $L(q)M^2(q^4) = 4L(q^4)M^2(q^4) + \frac{1}{87296}M(q)N(q) + \frac{255}{87296}M(q^2)N(q^2) - \frac{1024}{341}M(q^4)N(q^4) - \frac{23805}{992}C(q) - \frac{28125}{62}D(q) - \frac{39240}{31}E(q),$
- (vi) $M(q)N(q^4) = \frac{5}{21824}M(q)N(q) + \frac{315}{21824}M(q^2)N(q^2) + \frac{336}{341}M(q^4)N(q^4)$

CONVOLUTION SUMS

$$\begin{aligned}
 & + \frac{59535}{248}C(q) + \frac{187110}{31}D(q) + \frac{997920}{31}E(q), \\
 \text{(vii) } N(q)M(q^4) &= \frac{21}{5456}M(q)N(q) + \frac{315}{5456}M(q^2)N(q^2) + \frac{320}{341}M(q^4)N(q^4) \\
 & - \frac{31185}{62}C(q) - \frac{748440}{31}D(q) - \frac{7620480}{31}E(q), \\
 \text{(viii) } M^2(q)L(q^4) &= \frac{1}{4}L(q)M^2(q) - \frac{64}{341}M(q)N(q) + \frac{255}{1364}M(q^2)N(q^2) \\
 & + \frac{256}{341}M(q^4)N(q^4) + \frac{9810}{31}C(q) + \frac{900000}{31}D(q) + \frac{12188160}{31}E(q).
 \end{aligned}$$

Appealing to (2.1), (2.2), (2.3), (2.5), (2.9), (2.11), (6.1), (6.2), (6.3) and Theorem 6.1, we obtain the next result.

THEOREM 6.2.

$$\begin{aligned}
 \text{(i) } T_{7,1}(n) &= \frac{1}{14880}\sigma_9(n) + \frac{17}{744}\sigma_9(n/2) + \frac{(2-3n)}{48}\sigma_7(n/2) - \frac{1}{480}\sigma(n) \\
 & + \frac{1}{496}c(n) + \frac{2}{31}d(n), \\
 \text{(ii) } T_{5,3}(n) &= \frac{1}{31248}\sigma_9(n) + \frac{1}{465}\sigma_9(n/2) - \frac{1}{240}\sigma_5(n/2) + \frac{1}{504}\sigma_3(n) \\
 & - \frac{1}{496}c(n) - \frac{2}{31}d(n), \\
 \text{(iii) } T_{3,5}(n) &= \frac{1}{7440}\sigma_9(n) + \frac{4}{1953}\sigma_9(n/2) - \frac{1}{240}\sigma_5(n) + \frac{1}{504}\sigma_3(n/2) \\
 & + \frac{1}{248}c(n) + \frac{4}{31}d(n), \\
 \text{(iv) } T_{1,7}(n) &= \frac{17}{2976}\sigma_9(n) + \frac{8}{465}\sigma_9(n/2) + \frac{(4-3n)}{96}\sigma_7(n) - \frac{1}{480}\sigma(n/2) \\
 & - \frac{1}{62}c(n) - \frac{16}{31}d(n), \\
 \text{(v) } U_{7,1}(n) &= \frac{1}{3809280}\sigma_9(n) + \frac{17}{253952}\sigma_9(n/2) + \frac{17}{744}\sigma_9(n/4) + \frac{(2-3n)}{48}\sigma_7(n/4) \\
 & - \frac{1}{480}\sigma(n) + \frac{529}{253952}c(n) + \frac{625}{15872}d(n) + \frac{109}{992}e(n), \\
 \text{(vi) } U_{5,3}(n) &= \frac{1}{1999872}\sigma_9(n) + \frac{1}{31744}\sigma_9(n/2) + \frac{1}{465}\sigma_9(n/4) - \frac{1}{240}\sigma_5(n/4) \\
 & + \frac{1}{504}\sigma_3(n) - \frac{63}{31744}c(n) - \frac{99}{1984}d(n) - \frac{33}{124}e(n),
 \end{aligned}$$

$$\begin{aligned}
\text{(vii)} \quad U_{3,5}(n) &= \frac{1}{119040}\sigma_9(n) + \frac{1}{7936}\sigma_9(n/2) + \frac{4}{1953}\sigma_9(n/4) - \frac{1}{240}\sigma_5(n) \\
&\quad + \frac{1}{504}\sigma_3(n/4) + \frac{33}{7936}c(n) + \frac{99}{496}d(n) + \frac{63}{31}e(n), \\
\text{(viii)} \quad U_{1,7}(n) &= \frac{17}{11904}\sigma_9(n) + \frac{17}{3968}\sigma_9(n/2) + \frac{8}{465}\sigma_9(n/4) + \frac{(8-3n)}{192}\sigma_7(n) \\
&\quad - \frac{1}{480}\sigma(n/4) - \frac{109}{3968}c(n) - \frac{625}{248}d(n) - \frac{1058}{31}e(n).
\end{aligned}$$

7. Evaluation of $T_{e,f}(n)$ and $U_{e,f}(n)$ for $e + f = 10$

We first determine $L(q)M(q^2)N(q^2)$, $M(q)M^2(q^2)$, $N(q)N(q^2)$, $M^2(q)M(q^2)$, $M(q)N(q)L(q^2)$, $L(q)M(q^4)N(q^4)$, $M(q)M^2(q^4)$, $N(q)N(q^4)$, $M^2(q)M(q^4)$ and $M(q)N(q)L(q^4)$. In order to do this we introduce the constants $f(n)$ ($n \in \mathbb{N}$) defined by

$$(7.1) \quad F(q) = \sum_{n=1}^{\infty} f(n)q^n = \left(\frac{\Delta(q^4)^4}{\Delta(q)} \right)^{1/3} = \frac{x^5 w^{12}}{2^{20}}.$$

The first twenty values of $f(n)$ are given in the following table.

n	$f(n)$	n	$f(n)$	n	$f(n)$	n	$f(n)$
1	0	6	8	11	6296	16	388608
2	0	7	44	12	16384	17	756822
3	0	8	192	13	39569	18	1419200
4	0	9	694	14	89424	19	2572328
5	1	10	2208	15	191028	20	4521984

Similarly to the proof of Theorem 3.1, we obtain the following result.

THEOREM 7.1.

$$\begin{aligned}
\text{(i)} \quad L(q)M(q^2)N(q^2) &= 2L(q^2)M(q^2)N(q^2) - \frac{7}{520}M(q)^3 - \frac{56}{65}M(q^2)^3 \\
&\quad + \frac{31}{2184}N(q)^2 - \frac{38}{273}N(q^2)^2, \\
\text{(ii)} \quad M(q)M^2(q^2) &= \frac{11}{78}M(q)^3 + \frac{196}{39}M(q^2)^3 - \frac{25}{182}N(q)^2 - \frac{1100}{273}N(q^2)^2, \\
\text{(iii)} \quad N(q)N(q^2) &= -\frac{147}{520}M(q)^3 - \frac{1176}{65}M(q^2)^3 + \frac{31}{104}N(q)^2 + \frac{248}{13}N(q^2)^2,
\end{aligned}$$

$$(iv) \quad M^2(q)M(q^2) = \frac{49}{156}M(q)^3 + \frac{1408}{39}M(q^2)^3 - \frac{275}{1092}N(q)^2 - \frac{3200}{91}N(q^2)^2,$$

$$(v) \quad M(q)N(q)L(q^2) = \frac{1}{2}L(q)M(q)N(q) - \frac{14}{65}M(q)^3 - \frac{896}{65}M(q^2)^3 \\ - \frac{19}{546}N(q)^2 + \frac{3968}{273}N(q^2)^2,$$

$$(vi) \quad L(q)M(q^4)N(q^4) = 4L(q^4)M(q^4)N(q^4) - \frac{683}{49920}M(q)^3 - \frac{1727}{4160}M(q^2)^3 \\ - \frac{22138}{195}M(q^4)^3 + \frac{327}{23296}N(q)^2 + \frac{605}{1456}N(q^2)^2 \\ + \frac{10058}{91}N(q^4)^2 - 1520640F(q),$$

$$(vii) \quad M(q)M^2(q^4) = \frac{57}{416}M(q)^3 + \frac{515}{104}M(q^2)^3 + \frac{13932}{13}M(q^4)^3 \\ - \frac{175}{1248}N(q)^2 - \frac{450}{91}N(q^2)^2 - \frac{292300}{273}N(q^4)^2 \\ + 13824000F(q),$$

$$(viii) \quad N(q)N(q^4) = -\frac{4851}{16640}M(q)^3 - \frac{69237}{4160}M(q^2)^3 - \frac{77616}{65}M(q^4)^3 \\ + \frac{971}{3328}N(q)^2 + \frac{3465}{208}N(q^2)^2 + \frac{15536}{13}N(q^4)^2,$$

$$(ix) \quad M^2(q)M(q^4) = \frac{129}{416}M(q)^3 + \frac{3765}{104}M(q^2)^3 - \frac{89664}{13}M(q^4)^3 - \frac{2225}{8736}N(q)^2 \\ - \frac{13175}{364}N(q^2)^2 + \frac{1883200}{273}N(q^4)^2 - 221184000F(q),$$

$$(x) \quad M(q)N(q)L(q^4) = \frac{1}{4}L(q)M(q)N(q) - \frac{853}{3120}M(q)^3 - \frac{3971}{130}M(q^2)^3 \\ + \frac{2651392}{195}M(q^4)^3 - \frac{1}{208}N(q)^2 + \frac{2805}{91}N(q^2)^2 \\ - \frac{1237248}{91}N(q^4)^2 + 389283840F(q).$$

Appealing to (2.1), (2.2), (2.3), (2.6), (2.7), (2.11), (2.12), (7.1) and Theorem 7.1, we obtain the following result.

THEOREM 7.2.

- (i) $T_{9,1}(n) = \frac{1}{91212}\sigma_{11}(n) + \frac{31}{2073}\sigma_{11}(n/2) + \frac{(5-6n)}{120}\sigma_9(n/2) + \frac{1}{264}\sigma(n)$
 $- \frac{21}{5528}\tau(n) - \frac{282}{3455}\tau(n/2),$
- (ii) $T_{7,3}(n) = \frac{1}{331680}\sigma_{11}(n) + \frac{17}{20730}\sigma_{11}(n/2) - \frac{1}{240}\sigma_7(n/2) - \frac{1}{480}\sigma_3(n)$
 $+ \frac{23}{11056}\tau(n) + \frac{91}{1382}\tau(n/2),$
- (iii) $T_{5,5}(n) = \frac{1}{174132}\sigma_{11}(n) + \frac{16}{43533}\sigma_{11}(n/2) + \frac{1}{504}\sigma_5(n) + \frac{1}{504}\sigma_5(n/2)$
 $- \frac{11}{5528}\tau(n) - \frac{88}{691}\tau(n/2),$
- (iv) $T_{3,7}(n) = \frac{17}{331680}\sigma_{11}(n) + \frac{8}{10365}\sigma_{11}(n/2) - \frac{1}{240}\sigma_7(n) - \frac{1}{480}\sigma_3(n/2)$
 $+ \frac{91}{22112}\tau(n) + \frac{368}{691}\tau(n/2),$
- (v) $T_{1,9}(n) = \frac{31}{8292}\sigma_{11}(n) + \frac{256}{22803}\sigma_{11}(n/2) + \frac{(5-3n)}{120}\sigma_9(n) + \frac{1}{264}\sigma(n/2)$
 $- \frac{141}{6910}\tau(n) - \frac{2688}{691}\tau(n/2),$
- (vi) $U_{9,1}(n) = \frac{31}{5837568}\sigma_{11}(n) + \frac{1}{176896}\sigma_{11}(n/2) + \frac{31}{2073}\sigma_{11}(n/4)$
 $+ \frac{(5-6n)}{120}\sigma_9(n/4) + \frac{1}{264}\sigma(n),$
 $- \frac{671}{176896}\tau(n) - \frac{2505}{22112}\tau(n/2) - \frac{105314}{3455}\tau(n/4) - 240f(n),$
- (vii) $U_{7,3}(n) = -\frac{7}{2653440}\sigma_{11}(n) + \frac{1}{17689}\sigma_{11}(n/2) + \frac{17}{20730}\sigma_{11}(n/4)$
 $- \frac{1}{240}\sigma_7(n/4) - \frac{1}{480}\sigma_3(n)$
 $+ \frac{369}{176896}\tau(n) + \frac{1641}{22112}\tau(n/2) + \frac{22203}{1382}\tau(n/4) + 120f(n),$
- (viii) $U_{5,5}(n) = \frac{1}{11144448}\sigma_{11}(n) + \frac{1}{176896}\sigma_{11}(n/2) + \frac{16}{43533}\sigma_{11}(n/4)$
 $+ \frac{1}{504}\sigma_5(n) + \frac{1}{504}\sigma_5(n/4)$
 $- \frac{351}{176896}\tau(n) - \frac{2505}{22112}\tau(n/2) - \frac{5616}{691}\tau(n/4),$

CONVOLUTION SUMS

$$(ix) \ U_{3,7}(n) = \frac{121}{2653440}\sigma_{11}(n) + \frac{1}{176896}\sigma_{11}(n/2) + \frac{8}{10365}\sigma_{11}(n/4) \\ - \frac{1}{240}\sigma_7(n) - \frac{1}{480}\sigma_3(n/4) \\ + \frac{729}{176896}\tau(n) + \frac{6003}{11056}\tau(n/2) - \frac{71496}{691}\tau(n/4) - 1920f(n),$$

$$(x) \ U_{1,9}(n) = -\frac{7}{16584}\sigma_{11}(n) + \frac{23}{5528}\sigma_{11}(n/2) + \frac{256}{22803}\sigma_{11}(n/4) \\ + \frac{(10-3n)}{240}\sigma_9(n) + \frac{1}{264}\sigma(n/4) \\ - \frac{1589}{55280}\tau(n) - \frac{5790}{691}\tau(n/2) + \frac{2562304}{691}\tau(n/4) + 61440f(n).$$

8. Sums of 12 triangular numbers

The triangular numbers are the nonnegative integers

$$T_k = \frac{1}{2}k(k+1), \quad k = 0, 1, 2, \dots,$$

so that

$$T_0 = 0, \quad T_1 = 1, \quad T_2 = 3, \quad T_3 = 6, \quad T_4 = 10, \quad T_5 = 15, \dots$$

We set

$$\Delta = \{T_k \mid k = 0, 1, 2, \dots\}.$$

For $m \in \mathbb{N}$ and $n \in \mathbb{N} \cup \{0\}$, we let

$$\delta_m(n) = \text{card}\{(t_1, t_2, \dots, t_m) \in \Delta \times \Delta \times \dots \times \Delta \mid n = t_1 + t_2 + \dots + t_m\}$$

so that $\delta_m(n)$ counts the number of representations of n as the sum of m triangular numbers. For $n \in \mathbb{N} \cup \{0\}$ we have the classical results

$$(8.1) \quad \delta_4(n) = \sigma(2n+1)$$

and

$$(8.2) \quad \delta_8(n) = \sigma_3(n+1) - \sigma_3\left(\frac{n+1}{2}\right),$$

see for example [4, Theorem 10, p. 259; Theorem 12, p. 265], [10, Theorem 3, p. 80; Theorem 5, p. 82]. We use these two formulae in conjunction with Theorem 4.2 to give a very simple proof of the following result of Ono, Robins and Wahl [10, Theorem 7, p. 85], which was originally proved using the theory of modular forms.

THEOREM 8.1. For $n \in \mathbb{N} \cup \{0\}$

$$\delta_{12}(n) = \frac{1}{256}(\sigma_5(2n+3) - a(2n+3)),$$

where $a(n)$ is defined in (4.1).

Proof. We have by (8.1) and (8.2)

$$\begin{aligned} \delta_{12}(n) &= \sum_{m=0}^n \delta_4(m)\delta_8(n-m) \\ &= \sum_{m=0}^n \sigma(2m+1) \left(\sigma_3(n-m+1) - \sigma_3\left(\frac{n-m+1}{2}\right) \right) \\ &= A - B, \end{aligned}$$

where

$$\begin{aligned} A &= \sum_{m=0}^n \sigma(2m+1)\sigma_3(n-m+1) \\ &= \sum_{k=1}^{n+1} \sigma_3(k)\sigma(2n+3-2k) \\ &= T_{3,1}(2n+3) \\ &= \frac{1}{240}(\sigma_5(2n+3) - \sigma(2n+3)), \end{aligned}$$

by Theorem 4.2(i), and

$$\begin{aligned} B &= \sum_{m=0}^n \sigma(2m+1)\sigma_3\left(\frac{n-m+1}{2}\right) \\ &= \sum_{k < \frac{2n+3}{4}} \sigma_3(k)\sigma(2n+3-4k) \\ &= U_{3,1}(2n+3) \\ &= -\frac{1}{240}\sigma(2n+3) + \frac{1}{3840}\sigma_5(2n+3) + \frac{1}{256}a(2n+3), \end{aligned}$$

by Theorem 4.2(iii). Hence

$$\begin{aligned} \delta_{12}(n) &= \frac{1}{240}\sigma_5(2n+3) - \frac{1}{3840}\sigma_5(2n+3) - \frac{1}{256}a(2n+3) \\ &= \frac{1}{256}(\sigma_5(2n+3) - a(2n+3)) \end{aligned}$$

as asserted. \square

The authors would like to thank Mathieu Lemire who independently checked the values in the tables for them.

CONVOLUTION SUMS

References

- [1] B. C. Berndt, *Ramanujan's Notebooks, Part II*, Springer-Verlag, New York, 1989.
- [2] B. C. Berndt, *Ramanujan's Notebooks, Part III*, Springer-Verlag, New York, 1991.
- [3] J. W. L. Glaisher, *On the numbers of representations of a number as a sum of $2r$ squares, where $2r$ does not exceed eighteen*, Proc. London Math. Soc. **5** (1907), 479-490.
- [4] J. G. Huard, Z. M. Ou, B. K. Spearman and K. S. Williams, *Elementary evaluation of certain convolution sums involving divisor functions*, Number Theory for the Millennium II, edited by M. A. Bennett, B. C. Berndt, N. Boston, H. G. Diamond, A. J. Hildebrand, and W. Philipp, A. K. Peters, Natick, Massachusetts, 2002, pp. 229-274.
- [5] J. G. Huard and K. S. Williams, *Sums of sixteen squares*, Far East J. Math. Sci. **7** (2002), 147-164.
- [6] D. B. Lahiri, *On Ramanujan's function $\tau(n)$ and the divisor function $\sigma_k(n)$ -I*, Bull. Calcutta Math. Soc. **38** (1946), 193-206.
- [7] D. B. Lahiri, *On Ramanujan's function $\tau(n)$ and the divisor function $\sigma_k(n)$, Part II*, Bull. Calcutta Math. Soc. **39** (1947), 33-52.
- [8] G. Melfi, *Some problems in elementary number theory and modular forms*, Ph.D. Thesis, University of Pisa, 1997.
- [9] G. Melfi, *On some modular identities*, in Number Theory, Diophantine, Computational and Algebraic Aspects: Proceedings of the International Conference held in Eger, Hungary. Walter de Gruyter and Co. (1998), 371-382.
- [10] K. Ono, S. Robins and P. T. Wahl, *On the representation of integers as sums of triangular numbers*, Aequationes Math. **50** (1995), 73-94.
- [11] S. Ramanujan, *On certain arithmetical functions*, Trans. Cambridge Philos. Soc. **22** (1916), 159-184.
- [12] G. N. Watson, *A table of Ramanujan's function $\tau(n)$* , Proc. London Math. Soc. (2) **51** (1948), 1-13.

Nathalie Cheng and Kenneth S. Williams
Centre for Research in Algebra and Number Theory
School of Mathematics and Statistics
Carleton University
Ottawa, Ontario K1S 5B6
Canada
E-mail: nathaliecheng@yahoo.ca
williams@math.carleton.ca